arXiv:1511.02153v1 [math.CA] 6 Nov 2015

Complex-valued fractional derivatives on time scales*

Benaoumeur Bayour and Delfim F. M. Torres

Abstract We introduce a notion of fractional (noninteger order) derivative on an arbitrary nonempty closed subset of the real numbers (on a time scale). Main properties of the new operator are proved and several illustrative examples given.

Key words: fractional calculus; calculus on time scales; complex-valued operator; Hilger derivative of noninteger order.

2010 Mathematics Subject Classification: 26A33; 26E70.

1 Introduction

The study of fractional (noninteger) order derivatives on discrete, continuous and, more generally, on an arbitrary nonempty closed set (i.e., a time scale) is a well-known subject under strong current development. The subject is very rich and several different definitions and approaches are available, either in discrete [1], continuous [11], and time-scale settings [2]. In continuous time, i.e., for the time scale $\mathbb{T} = \mathbb{R}$, several definitions are based on the classical Euler Gamma function Γ. For the time scale $\mathbb{T} = \mathbb{Z}$, the Gamma function is nothing else than the factorial, while for the q-scale one has the q-Gamma function Γ_q [9]. For the definition of Gamma function on an arbitrary time scale $\mathbb{T}$ see [6]. Similarly to [2, 3], here we introduce a

Benaoumeur Bayour
University of Chlef, B.P. 151, Hay Es-salem Chlef, Algeria, e-mail: b.benaoumeur@gmail.com

Delfim F. M. Torres ✉
University of Aveiro, Center for Research and Development in Mathematics and Applications (CIDMA), Department of Mathematics, 3810-193 Aveiro, Portugal, e-mail: delfim@ua.pt

* This is a preprint of a paper whose final and definite form will appear in *Springer Proceedings in Mathematics & Statistics*, ISSN: 2194-1009. Accepted for publication 06/Nov/2015.

new notion of fractional derivative on an arbitrary time scale $\mathbb{T}$ that does not involve Gamma functions. Our approach is, however, different from the ones available in the literature [2, 3, 4, 5]. In particular, while in [2, 3, 4, 5] the fractional derivative at a point is always a real number, here, in contrast, the fractional derivative at a point is, in general, a complex number. For example, the derivative of order $\alpha \in (0,1]$ of the square function t^2 is always given by $t^\alpha + (\sigma(t))^\alpha$, where $\sigma(t)$ is the forward jump operator of the time scale, which is in general a complex number (e.g., for $\alpha = 1/2$ and $t < 0$) and a generalization of the Hilger derivative $(t^2)^\Delta = t + \sigma(t)$.

The text is organized as follows. In Section 2 we recall the notion of Hilger/delta derivative. Our complex-valued fractional derivative on time scales is introduced in Section 3, where its main properties are proved and several examples given.

2 Preliminaries

A time scale $\mathbb{T}$ is an arbitrary nonempty closed subset of the real numbers $\mathbb{R}$. For $t \in \mathbb{T}$, we define the forward jump operator $\sigma : \mathbb{T} \to \mathbb{T}$ by $\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}$ and the backward jump operator $\rho : \mathbb{T} \to \mathbb{T}$ is defined by $\rho(t) := \sup\{s \in \mathbb{T} : s < t\}$. Then, one defines the graininess function $\mu : \mathbb{T} \to [0, +\infty[$ by $\mu(t) = \sigma(t) - t$.

If $\sigma(t) > t$, then we say that t is right-scattered; if $\rho(t) < t$, then t is left-scattered. Moreover, if $t < \sup \mathbb{T}$ and $\sigma(t) = t$, then t is called right-dense; if $t > \inf \mathbb{T}$ and $\rho(t) = t$, then t is called left-dense. If $\mathbb{T}$ has a left-scattered maximum m, then we define $\mathbb{T}^\kappa = \mathbb{T} \setminus \{m\}$; otherwise $\mathbb{T}^\kappa = \mathbb{T}$. If $f : \mathbb{T} \to \mathbb{R}$, then $f^\sigma : \mathbb{T} \to \mathbb{R}$ is given by $f^\sigma(t) = f(\sigma(t))$ for all $t \in \mathbb{T}$.

Definition 1 (The Hilger derivative [8]). Let $f : \mathbb{T} \to \mathbb{R}$ and $t \in \mathbb{T}$. We define $f^\Delta(t)$ to be the number (provided it exists) with the property that given any $\varepsilon > 0$ there is a neighborhood U of t (i.e., $U = (t - \delta, t + \delta) \cap \mathbb{T}$ for some $\delta > 0$) such that

$$|[f(\sigma(t)) - f(s)] - f^\Delta(t)[\sigma(t) - s]| \leq \varepsilon |\sigma(t) - s|$$

for all $s \in U$. We call $f^\Delta(t)$ the Hilger (or delta) derivative of f at t.

For more on the calculus on time scales we refer the reader to the books [7, 8].

3 Complex-valued fractional derivatives on time scales

Let $f : \mathbb{T} \to \mathbb{R}$ with $\mathbb{T}$ a given time scale. We introduce here a new definition of fractional (noninteger) delta derivative of order $\alpha \in (0,1]$ at a point $t \in \mathbb{T}^\kappa$.

Definition 2 (The delta fractional derivative of order α). Assume $f : \mathbb{T} \to \mathbb{R}$ with $\mathbb{T}$ a time scale. Let $t \in \mathbb{T}^\kappa$ and $\alpha \in (0,1]$. We define ${f^\Delta}^\alpha(t)$ to be the number (provided it exists) with the property that given any $\varepsilon > 0$ there is a neighborhood U of t (i.e., $U = (t - \delta, t + \delta) \cap \mathbb{T}$ for some $\delta > 0$) such that

$$\left|[f^{\alpha}(\sigma(t))-f^{\alpha}(s)]-f^{\Delta^{\alpha}}(t)[\sigma(t)^{\alpha}-s^{\alpha}]\right|\leq\varepsilon\,|\sigma(t)^{\alpha}-s^{\alpha}| \tag{1}$$

for all $s\in U$. We call $f^{\Delta^{\alpha}}(t)$ the delta derivative of order α of f at t or the delta fractional (noninteger order) derivative of f at t. Moreover, we say that f is delta differentiable of order α on $\mathbb{T}^{\kappa}$ provided $f^{\Delta^{\alpha}}(t)$ exists for all $t\in\mathbb{T}^{\kappa}$. Function $f^{\Delta^{\alpha}}:\mathbb{T}^{\kappa}\to\mathbb{C}$ is then called the delta derivative of order α of f on $\mathbb{T}^{\kappa}$.

Remark 1. In (1) we use f^{α} to denote the power α of f. It is clear that the new derivative coincides with the standard Hilger derivative in the integer order case $\alpha=1$. Differently from $\alpha=1$, in general $f^{\Delta^{\alpha}}(t)$ is a complex number.

Theorem 1. *Assume $f:\mathbb{T}\to\mathbb{R}$ with $\mathbb{T}$ a time scale. Let $t\in\mathbb{T}^{\kappa}$ and $\alpha\in\mathbb{R}$. Then the following proprieties hold:*

1. *If f is continuous at t and t is right-scattered, then f is delta differentiable of order α at t with*
$$f^{\Delta^{\alpha}}(t)=\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(t)}{\sigma^{\alpha}(t)-t^{\alpha}}. \tag{2}$$
2. *If t is right-dense, then f is delta differentiable of order α at t if only if the limit*
$$\lim_{s\to t}\frac{f^{\alpha}(t)-f^{\alpha}(s)}{t^{\alpha}-s^{\alpha}}$$
exists as a finite number. In this case
$$f^{\Delta^{\alpha}}(t)=\lim_{s\to t}\frac{f^{\alpha}(t)-f^{\alpha}(s)}{t^{\alpha}-s^{\alpha}}. \tag{3}$$
3. *If f is delta differentiable of order α at t, then*
$$f^{\alpha}(\sigma(t))=f^{\alpha}(t)+(\sigma(t)^{\alpha}-t^{\alpha})f^{\Delta^{\alpha}}(t).$$

Proof. 1. Assume f is continuous at t and t is right scattered. By continuity,

$$\lim_{s\to t}\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(s)}{\sigma^{\alpha}(t)-s^{\alpha}}=\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(t)}{\sigma^{\alpha}(t)-t^{\alpha}}.$$

Hence, given $\varepsilon>0$, there is a neighborhood U of t such that

$$\left|\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(s)}{\sigma^{\alpha}(t)-s^{\alpha}}-\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(t)}{\sigma^{\alpha}(t)-t^{\alpha}}\right|\leq\varepsilon$$

for all $s\in U$. It follows that

$$\left|f^{\alpha}(\sigma(t))-f^{\alpha}(s)-\frac{f^{\alpha}(\sigma(t))-f^{\alpha}(t)}{\sigma^{\alpha}(t)-t^{\alpha}}[\sigma^{\alpha}(t)-s^{\alpha}]\right|\leq\varepsilon\,|\sigma^{\alpha}(t)-s^{\alpha}|$$

for all $s\in U$. Hence, we get the desired result (2).

2. Assume f is differentiable at t and t is right-dense. Let $\varepsilon > 0$ be given. Since f is differentiable at t, there is a neighborhood U of t such that

$$| [f^{\alpha}(\sigma(t)) - f^{\alpha}(t)] - f^{\Delta^{\alpha}}(t)[\sigma^{\alpha}(t) - s^{\alpha}] | \leq \varepsilon \, | \, \sigma^{\alpha}(t) - s^{\alpha} \, |$$

for all $s \in U$. Since $\sigma(t) = t$, we have that

$$| [f^{\alpha}(\sigma(t)) - f^{\alpha}(t)] - f^{\Delta^{\alpha}}(t)[t^{\alpha} - s^{\alpha}] | \leq \varepsilon \, | \, \sigma^{\alpha}(t) - s^{\alpha} \, |$$

for all $s \in U$. It follows that $\left| \frac{f^{\alpha}(t) - f^{\alpha}(s)}{t^{\alpha} - s^{\alpha}} - f^{\Delta^{\alpha}}(t) \right| \leq \varepsilon$ for all $s \in U$, $s \neq t$, and we get the desired equality (3). Assume $\lim_{s \to t} \frac{f^{\alpha}(t) - f^{\alpha}(s)}{t^{\alpha} - s^{\alpha}}$ exists and is equal to X and $\sigma(t) = t$. Let $\varepsilon > 0$. Then there is a neighborhood U of t such that

$$\left| \frac{f^{\alpha}(\sigma(t)) - f^{\alpha}(s)}{t^{\alpha} - s^{\alpha}} - X \right| \leq \varepsilon$$

for all $s \in U$. Because $| \, f^{\alpha}(\sigma(t)) - f^{\alpha}(s) - X(t^{\alpha} - s^{\alpha}) \, | \leq \varepsilon |t^{\alpha} - s^{\alpha}|$ for all $s \in U$,

$$f^{\Delta^{\alpha}}(t) = X = \lim_{s \to t} \frac{f^{\alpha}(t) - f^{\alpha}(s)}{t^{\alpha} - s^{\alpha}}.$$

3. If $\sigma(t) = t$, then $\sigma^{\alpha}(t) - t^{\alpha} = 0$ and

$$f^{\alpha}(\sigma(t)) = f^{\alpha}(t) = f^{\alpha}(t) + (\sigma^{\alpha}(t) - t^{\alpha}) f^{\Delta^{\alpha}}(t).$$

On the other hand, if $\sigma(t) > t$, then by item 1

$$\begin{aligned} f^{\alpha}(\sigma(t)) &= f^{\alpha}(t) + (\sigma^{\alpha}(t) - t^{\alpha}) \frac{f^{\alpha}(\sigma(t)) - f^{\alpha}(t)}{\sigma(t)^{\alpha} - t^{\alpha}} \\ &= f^{\alpha}(t) + (\sigma^{\alpha}(t) - t^{\alpha}) f^{\Delta^{\alpha}}(t) \end{aligned}$$

and the proof is complete.

Example 1. If $\mathbb{T} = \mathbb{R}$, then (3) yields that $f : \mathbb{R} \to \mathbb{R}$ is delta differentiable of order α at $t \in \mathbb{R}$ if and only if $f^{\Delta^{\alpha}}(t) = \lim_{s \to t} \frac{f^{\alpha}(t) - f^{\alpha}(s)}{t^{\alpha} - s^{\alpha}}$ exists, i.e., if and only if f is fractional differentiable at t. In this case we get the derivative $f^{(\alpha)}$ of [10].

Example 2. If $\mathbb{T} = \mathbb{Z}$, then item 1 of Theorem 1 yields that $f : \mathbb{Z} \to \mathbb{R}$ is delta-differentiable of order α at $t \in \mathbb{Z}$ with

$$f^{\Delta^{\alpha}}(t) = \frac{f^{\alpha}(\sigma(t)) - f^{\alpha}(t)}{\sigma^{\alpha}(t) - t^{\alpha}} = \frac{f^{\alpha}(t+1) - f^{\alpha}(t)}{(t+1)^{\alpha} - t^{\alpha}}.$$

Example 3. If $f : \mathbb{T} \to \mathbb{R}$ is defined by $f(t) \equiv \lambda \in \mathbb{R}$, then $f^{\Delta^{\alpha}}(t) \equiv 0$. Indeed, if t is right-scattered, then by item 1 of Theorem 1 $f^{\Delta^{\alpha}}(t) = \frac{f^{\alpha}(\sigma(t)) - f^{\alpha}(t)}{\sigma^{\alpha}(t) - t^{\alpha}} = \frac{\lambda^{\alpha} - \lambda^{\alpha}}{\sigma^{\alpha}(t) - t^{\alpha}} = 0$; if t is right-dense, then by (3) we get $f^{\Delta^{\alpha}}(t) = \lim_{s \to t} \frac{\lambda^{\alpha} - \lambda^{\alpha}}{t^{\alpha} - s^{\alpha}} = 0$.

Example 4. If $f:\mathbb{T}\to\mathbb{R}$, $t\mapsto t$, then $f^{\Delta^\alpha}\equiv 1$ because if $\sigma(t)>t$ (i.e., t is right-scattered), then $f^{\Delta^\alpha}(t)=\frac{f^\alpha(\sigma(t))-f^\alpha(t)}{\sigma^\alpha(t)-t^\alpha}=\frac{\sigma^\alpha(t)-t^\alpha}{\sigma^\alpha(t)-t^\alpha}=1$; if $\sigma(t)=t$ (i.e., t is right-dense), then $f^{\Delta^\alpha}=\lim_{s\to t}\frac{f^\alpha(t)-f^\alpha(s)}{t^\alpha-s^\alpha}=\frac{t^\alpha-s^\alpha}{t^\alpha-s^\alpha}=1$.

Example 5. Let $g:\mathbb{T}\to\mathbb{R}$, $t\mapsto\frac{1}{t}$. We have $g^{\Delta^\alpha}(t)=-\frac{1}{(t\sigma(t))^\alpha}$. Indeed, if $\sigma(t)=t$, then $g^{\Delta^\alpha}(t)=-\frac{1}{t^{2\alpha}}$; if $\sigma(t)>t$, then

$$g^{\Delta^\alpha}(t)=\frac{g^\alpha(\sigma(t))-g^\alpha(t)}{\sigma^\alpha(t)-t^\alpha}=\frac{\left(\frac{1}{\sigma(t)}\right)^\alpha-\left(\frac{1}{t}\right)^\alpha}{\sigma^\alpha(t)-t^\alpha}=\frac{\frac{t^\alpha-\sigma^\alpha(t)}{t^\alpha\sigma^\alpha(t)}}{t^\alpha-\sigma^\alpha(t)}=-\frac{1}{t^\alpha\sigma^\alpha(t)}.$$

Example 6. Let $h:\mathbb{T}\to\mathbb{R}$, $t\mapsto t^2$. We have $h^{\Delta^\alpha}(t)=\sigma^\alpha(t)+t^\alpha$. Indeed, if t is right-dense, then $h^{\Delta^\alpha}(t)=\lim_{s\to t}\frac{t^{2\alpha}-s^{2\alpha}}{t^\alpha-s^\alpha}=2t^\alpha$; if t is right-scattered, then

$$h^{\Delta^\alpha}(t)=\frac{h^\alpha(\sigma(t))-h^\alpha(t)}{\sigma^\alpha(t)-t^\alpha}=\frac{\sigma^{2\alpha}(t)-t^{2\alpha}}{\sigma^\alpha(t)-t^\alpha}=\sigma^\alpha(t)+t^\alpha.$$

Example 7. Consider the time scale $\mathbb{T}=h\mathbb{Z}$, $h>0$. Let f be the function defined by $f:h\mathbb{Z}\to\mathbb{R}$, $t\mapsto(t-c)^2$, $c\in\mathbb{R}$. The fractional derivative of order α of f at t is

$$\begin{aligned}f^{\Delta^\alpha}(t)&=\frac{f^\alpha(\sigma(t))-f^\alpha(t)}{\sigma^\alpha(t)-t^\alpha}=\frac{((\sigma(t)-c)^2)^\alpha-((t-c)^2)^\alpha}{\sigma^\alpha(t)-t^\alpha}\\&=\frac{(t+h-c)^{2\alpha}-(t-c)^{2\alpha}}{(t+h)^\alpha-t^\alpha}.\end{aligned}$$

Remark 2. Examples 5, 6 and 7 show that in general $f^{\Delta^\alpha}(t)$ is a complex number (for instance, choose $\alpha=\frac{1}{2}$ and $t<0$).

Theorem 2. *Assume $f,g:\mathbb{T}\to\mathbb{R}$ are continuous and delta differentiable of order α at $t\in\mathbb{T}^\kappa$. Then the following proprieties hold:*

1. *For any constant λ, function $\lambda f:\mathbb{T}\to\mathbb{R}$ is delta differentiable of order α at t with $(\lambda f)^{\Delta^\alpha}=\lambda^\alpha f^{\Delta^\alpha}$.*
2. *The product $fg:\mathbb{T}\to\mathbb{R}$ is delta differentiable of order α at t with*

$$(fg)^{\Delta^\alpha}(t)=f^{\Delta^\alpha}(t)g^\alpha(t)+f^\alpha(\sigma(t))g^{\Delta^\alpha}(t)=f^{\Delta^\alpha}(t)g^\alpha(\sigma(t))+f^\alpha(t)g^{\Delta^\alpha}(t).$$

3. *If $f(t)f(\sigma(t))\neq 0$, then $\frac{1}{f}$ is delta differentiable of order α at t with*

$$\left(\frac{1}{f}\right)^{\Delta^\alpha}(t)=\frac{-f^{\Delta^\alpha}(t)}{f^\alpha(\sigma(t))f^\alpha(t)}.$$

4. *If $g(t)g(\sigma(t))\neq 0$, then $\frac{f}{g}$ is delta differentiable of order α at t with*

$$\left(\frac{f}{g}\right)^{\Delta^\alpha}(t)=\frac{f^{\Delta^\alpha}(t)g^\alpha(t)-f^\alpha(t)g^{\Delta^\alpha}(t)}{g^\alpha(\sigma(t))g^\alpha(t)}.$$

Proof. 1. Let $\varepsilon \in (0,1)$. Define $\varepsilon^* = \frac{\varepsilon}{|\lambda|^\alpha} \in (0,1)$. Then there exists a neighborhood U of t such that $|f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)| \leq \varepsilon^* |\sigma^\alpha(t) - s^\alpha|$ for all $s \in U$. It follows that

$$\begin{aligned}
&|(\lambda f)^\alpha(\sigma(t)) - (\lambda f)^\alpha(s) - \lambda f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)| \\
&\quad = |\lambda|^\alpha \mid f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha) \mid \\
&\quad \leq \varepsilon^* |\lambda|^\alpha |\sigma^\alpha(t) - s^\alpha| \leq \frac{\varepsilon}{|\lambda|^\alpha} |\lambda|^\alpha |\sigma^\alpha(t) - s^\alpha| = \varepsilon |\sigma^\alpha(t) - s^\alpha|
\end{aligned}$$

for all $s \in U$. Thus $(\lambda f)^{\Delta^\alpha}(t) = \lambda^\alpha f^{\Delta^\alpha}(t)$ holds.

2. Let $\varepsilon \in (0,1)$. Define $\varepsilon^* = \varepsilon[1 + |f^\alpha(t)| + |g^\alpha(\sigma(t))| + |g^{\Delta^\alpha}(\sigma(t))|]^{-1}$. Then $\varepsilon^* \in (0,1)$ and there exist neighborhoods U_1, U_2 and U_3 of t such that

$$|f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)| \leq \varepsilon^* |\sigma^\alpha(t) - s^\alpha|$$

for all $s \in U_1$, $|g^\alpha(\sigma(t)) - g^\alpha(s) - g^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)| \leq \varepsilon^* |\sigma^\alpha(t) - s^\alpha|$ for all $s \in U_2$ and such as f is continuous. Then $|f(t) - f(s)| \leq \varepsilon^*$ for all $s \in U_3$. Define $U = U_1 \cap U_2 \cap U_3$ and let $s \in U$. It follows that

$$\begin{aligned}
&|(fg)^\alpha(\sigma(t)) - (fg)^\alpha(s) - [g^{\Delta^\alpha}(t) f^\alpha(t) + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)][\sigma^\alpha(t) - s^\alpha]| \\
&= |[f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)](g^\alpha(\sigma(t))) + g^\alpha(\sigma(t)) f^\alpha(s) \\
&\quad + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha) - f^\alpha(s) g^\alpha(s) \\
&\quad - [g^{\Delta^\alpha}(t) f^\alpha(t) + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)][\sigma^\alpha(t) - s^\alpha]| \\
&= |[f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)](g^\alpha(\sigma(t))) \\
&\quad + [g^\alpha(\sigma(t)) - g^\alpha(s) - g^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)](f^\alpha(t)) \\
&\quad + [g^\alpha(\sigma(t)) - g^\alpha(s) - g^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)](f^\alpha(s) - f^\alpha(t)) + f^\alpha(s) g^\alpha(s) \\
&\quad + g^{\Delta^\alpha}(t) f^\alpha(s)(\sigma^\alpha(t) - s^\alpha) + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha) - g^\alpha(s) f^\alpha(s) \\
&\quad + g^{\Delta^\alpha}(t) f^\alpha(s)(\sigma^\alpha(t) - s^\alpha) + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha) - f^\alpha(s) g^\alpha(s) \\
&\quad - [g^{\Delta^\alpha}(t) f^\alpha(t) + g^\alpha(\sigma(t)) f^{\Delta^\alpha}(t)][\sigma^\alpha(t) - s^\alpha]| \\
&\leq |f^\alpha(\sigma(t)) - f^\alpha(s) - f^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)||(g^\alpha(\sigma(t)))| \\
&\quad + |g^\alpha(\sigma(t)) - g^\alpha(s) - g^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)||(f^\alpha(t))| \\
&\quad + |g^\alpha(\sigma(t)) - g^\alpha(s) - g^{\Delta^\alpha}(t)(\sigma^\alpha(t) - s^\alpha)||f^\alpha(s) - f^\alpha(t)| \\
&\quad + |g^{\Delta^\alpha}(t)||f^\alpha(t) - f^\alpha(s)||\sigma^\alpha(t) - s^\alpha| \\
&= \varepsilon^* |(g^\alpha(\sigma(t)))||\sigma^\alpha(t) - s^\alpha| \\
&\quad + \varepsilon^* |(f^\alpha(t))||\sigma^\alpha(t) - s^\alpha| + \varepsilon^* |\sigma^\alpha(t) - s^\alpha| \varepsilon^* + \varepsilon^* |g^{\Delta^\alpha}(t)||\sigma^\alpha(t) - s^\alpha| \\
&\leq \varepsilon^* |\sigma^\alpha(t) - s^\alpha| (\varepsilon^* + |(f^\alpha(t))| + |g^{\Delta^\alpha}(t)| + |g^{\Delta^\alpha}(t)|) \\
&\leq \varepsilon^* |\sigma^\alpha(t) - s^\alpha| (1 + |(f^\alpha(t))| + |g^{\Delta^\alpha}(t)| + |g^{\Delta^\alpha}(t)|) = \varepsilon |\sigma^\alpha(t) - s^\alpha|.
\end{aligned}$$

Thus $(fg)^{\Delta^\alpha}(t) = f^\alpha(t)g^{\Delta^\alpha}(t) + f^{\Delta^\alpha}(t)g^\alpha(\sigma(t))$ holds at t. The other product rule follows from this last equality by interchanging functions f and g.

3. We use the delta derivative of a constant (Example 3). Since $\left(f\cdot\frac{1}{f}\right)^{\Delta^\alpha}(t) = 0$, it follows from item 2 that $\left(\frac{1}{f}\right)^{\Delta^\alpha}(t)f^\alpha(\sigma(t)) + f^{\Delta^\alpha}(t)\frac{1}{f^\alpha(t)} = 0$. Because we are assuming $f(t)f(\sigma(t)) \neq 0$, one has $\left(\frac{1}{f}\right)^{\Delta^\alpha}(t) = \frac{-f^{\Delta^\alpha}(t)}{f^\alpha(\sigma(t))f^\alpha(t)}$.
4. For the quotient formula we use items 2 and 3 to compute

$$\begin{aligned}\left(\frac{f}{g}\right)^{\Delta^\alpha}(t) &= \left(f\cdot\frac{1}{g}\right)^{\Delta^\alpha}(t) = f^\alpha(t)\left(\frac{1}{g}\right)^{\Delta^\alpha}(t) + f^{\Delta^\alpha}(t)\frac{1}{g^\alpha(\sigma(t))}\\ &= -f^\alpha(t)\frac{g^{\Delta^\alpha}(t)}{g^\alpha(\sigma(t))g^\alpha(t)} + f^{\Delta^\alpha}(t)\frac{1}{g^\alpha(\sigma(t))}\\ &= \frac{f^{\Delta^\alpha}(t)g^\alpha(t) - f^\alpha(t)g^{\Delta^\alpha}(t)}{g^\alpha(\sigma(t))g^\alpha(t)}.\end{aligned}$$

This concludes the proof.

Remark 3. The delta derivative of order α of the sum $f+g : \mathbb{T}\to\mathbb{R}$ does not satisfy the usual property, that is, in general $(f+g)^{\Delta^\alpha}(t) \neq (f)^{\Delta^\alpha}(t) + (g)^{\Delta^\alpha}(t)$. For instance, let $\mathbb{T}$ be an arbitrary time scale and f,g be functions defined by $f : \mathbb{T}\to\mathbb{R}$, $t\mapsto t$, and $g : \mathbb{T}\to\mathbb{R}$, $t\mapsto 2t$. One can easily find that $(f+g)^{\Delta^\alpha}(t) = \sqrt{3} \neq f^{\Delta^\alpha}(t) + g^{\Delta^\alpha}(t) = 1+\sqrt{2}$.

Proposition 1. *Let $\alpha\in\mathbb{R}$ and $m\in\mathbb{N}$, $m>1$. For g defined by $g(t)=t^m$ we have*

$$g^{\Delta^\alpha}(t) = \sum_{k=0}^{m-1}(t^\alpha)^{m-k-1}(\sigma^\alpha)^k(t). \tag{4}$$

Proof. We prove the formula by induction. If $m=2$, then $g(t)=t^2$ and from Example 6 we know that $g^{\Delta^\alpha}(t) = \sum_{k=0}^{1}(t^\alpha)^{1-k}(\sigma^\alpha)^k(t) = t^\alpha + \sigma^\alpha(t)$. Now assume

$$g^{\Delta^\alpha}(t) = \sum_{k=0}^{m-1}(t^\alpha)^{m-k-1}(\sigma^\alpha)^k(t)$$

holds for $g(t)=t^m$ and let $G(t) = t^{m+1} = t\cdot g(t)$. We use the product rule of Theorem 2 to obtain

$$\begin{aligned}G^{\Delta^\alpha}(t) &= g^\alpha(t) + \sigma^\alpha(t)g^{\Delta^\alpha}(t) = (t^\alpha)^m + \sigma^\alpha(t)\sum_{k=0}^{m-1}(t^\alpha)^{m-k-1}(\sigma^\alpha)^k(t)\\ &= (t^\alpha)^m + \sum_{k=0}^{m-1}(t^\alpha)^{m-k-1}(\sigma^\alpha)^{k+1}(t) = (t^\alpha)^m + \sum_{k=1}^{m-1}(t^\alpha)^{m-k}(\sigma^\alpha)^k(t)\\ &= \sum_{k=0}^{m}(t^\alpha)^{m-k}(\sigma^\alpha)^k(t).\end{aligned}$$

Hence, by mathematical induction, (4) holds.

Example 8. Choose $m = 3$ in Proposition 1. Then $\left(t^3\right)^{\Delta^\alpha} = t^{2\alpha} + (t\sigma(t))^\alpha + \sigma^{2\alpha}(t)$.

The notion of fractional derivative here introduced can be easily extended to any arbitrary real order α.

Definition 3. Let $\alpha > 0$ and $N \in \mathbb{N}_0$ be such that $N < \alpha \leq N+1$. Then we define $f^{\Delta^\alpha} = \left(f^{\Delta^N}\right)^{\Delta^{\alpha-N}}$, where f^{Δ^N} is the usual Hilger derivative of order N.

Acknowledgements This research is part of first author's Ph.D. project, which is carried out at Sidi Bel Abbes University, Algeria. It was initiated while Bayour was visiting the Department of Mathematics of University of Aveiro, Portugal, 2015. The hospitality of the host institution and the financial support of University of Chlef, Algeria, are here gratefully acknowledged. Torres was supported by Portuguese funds through CIDMA and FCT, within project UID/MAT/04106/2013.

References

1. F. M. Atıcı and M. Uyanik, Analysis of discrete fractional operators, Appl. Anal. Discrete Math. **9** (2015), no. 1, 139–149.
2. N. Benkhettou, A. M. C. Brito da Cruz and D. F. M. Torres, A fractional calculus on arbitrary time scales: fractional differentiation and fractional integration, Signal Process. **107** (2015), 230–237. arXiv:1405.2813
3. N. Benkhettou, A. M. C. Brito da Cruz and D. F. M. Torres, Nonsymmetric and symmetric fractional calculi on arbitrary nonempty closed sets, Mathematical Methods in the Applied Sciences, in press. DOI:10.1002/mma.3475 arXiv:1502.07277
4. N. Benkhettou, A. Hammoudi and D. F. M. Torres, Existence and uniqueness of solution for a fractional Riemann–Liouville initial value problem on time scales, J. King Saud Univ. Sci., in press. DOI:10.1016/j.jksus.2015.08.001 arXiv:1508.00754
5. N. Benkhettou, S. Hassani and D. F. M. Torres, A conformable fractional calculus on arbitrary time scales, J. King Saud Univ. Sci., in press. DOI:10.1016/j.jksus.2015.05.003 arXiv:1505.03134
6. M. Bohner and B. Karpuz, The gamma function on time scales, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. **20** (2013), no. 4, 507–522.
7. M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser Boston, Boston, MA, 2001.
8. M. Bohner and A. Peterson, *Advances in dynamic equations on time scales*, Birkhäuser Boston, Boston, MA, 2003.
9. V. Kac and P. Cheung, *Quantum calculus*, Universitext, Springer, New York, 2002.
10. A. Karci and A. Karadogan, Fractional order derivative and relationship between derivative and complex functions, Math. Sci. Appl. E-Notes **2** (2014), no. 1, 44–54.
11. G. S. Taverna and D. F. M. Torres, Generalized fractional operators for nonstandard Lagrangians, Math. Methods Appl. Sci. **38** (2015), no. 9, 1808–1812. arXiv:1404.6483